Topic: Other

Preference: Oral

Discrete Breathers in 2d SRR systems

M. Eleftheriou^{1*}, N. Lazarides², G. P. Tsironis³

 ¹ Technological Educational Institute of Crete at Rethymnon, E. Daskalaki, Perivolia, 74100 Rethymno, Crete, Greece and Department of Materials Science and Technology, University of Crete, P. O. Box 2208, Greece
² Department of Electrical Engineering, Technological Educational Institute of Crete, P. O. Box 140, Stavromenos, 71500, Heraklion, Crete, Greece and Department of Physics, University of Crete, P. O. Box 2208, 71003 Heraklion, Greece and Institute of Electronic Structure and Laser, FORTH, P. O. Box 1527, 71110 Heraklion, Greece
³ Department of Physics, University of Crete, P. O. Box 2208, 71003
Heraklion,Greece and Institute of Electronic Structure and Laser, FORTH, P. O. Box 1527, 71110 Heraklion, Greece
* Electronic Address: marel@physics.uoc.gr

We consider SRR (split ring resonators) arrays positioned in the 2D plane, with axial or planar geometry. Using these topologies we construct Discrete Breathers of various types such as Hamiltonian, dissipative, single site and multibreathers as well as surface DB modes, located in the corner or in the edges of the lattice. We show that the presence of these modes induce negative magnetic response in the system.

- N. Lazarides, M. Eleftheriou, and G. P. Tsironis, *Phys. Rev. Lett.* 97, 157406 (2006).
- [2] M. Eleftheriou, N. Lazarides and G. P. Tsironis, *Phys. Rev. E* 77, 036608 (2008).
- [3] N. Lazarides, G. P. Tsironis, Y. Kivshar, Phys. Rev. E 77, 065601 (2008).